

Efficiency analysis of double perturbed pairwise comparison matrices

Kristóf Ábele-Nagy^{1,2*} Sándor Bozóki^{1,2†} Örs Rebák[‡]

Abstract

Efficiency is a core concept of multi-objective optimization problems and multi-attribute decision making. In the case of pairwise comparison matrices a weight vector is called efficient if the approximations of the elements of the pairwise comparison matrix made by the ratios of the weights cannot be improved in any position without making it worse in some other position. A pairwise comparison matrix is called double perturbed if it can be made consistent by altering two elements and their reciprocals. The most frequently used weighting method, the eigenvector method is analyzed in the paper, and it is shown that it produces an efficient weight vector for double perturbed pairwise comparison matrices.

1 Introduction

Ranking alternatives, or picking the best alternative is a commonly investigated problem. The case of a single cardinal objective function to be maximized or minimized is long studied by various operations research disciplines. This is however often not feasible. Alternatives can be ranked by assigning a cardinal utility to them, or by setting up ordinal preference relations among them. In the case of a single criterion and a single decision maker, modelling the preferences is often possible through standard methods. If there are multiple, often contradicting criteria, this becomes significantly harder. A dominant alternative, which is the best with respect to all criteria, very rarely exists. Thus, when a decision making method is used to aid the decision of a decision maker, some form of compromise is needed. Modelling the preferences of the decision maker by ranking or weighting the criteria can accomplish such a compromise. It allows the “best” alternative to be chosen (or the possible alternatives to be ranked) with respect to the subjective preferences of the decision maker. Examples of multi-criteria decision problems range from “Which house to buy?” or “What should the company invest in?” to public tenders.

*E-mail: abele-nagy.kristof@sztaki.mta.hu

†E-mail: bozoki.sandor@sztaki.mta.hu

‡E-mail: rebakors@gmail.com

¹Laboratory on Engineering and Management Intelligence, Research Group of Operations Research and Decision Systems, Computer and Automation Research Institute, Hungarian Academy of Sciences; Mail: 1518 Budapest, P.O. Box 63, Hungary.

²Department of Operations Research and Actuarial Sciences, Corvinus University of Budapest, Hungary

When weighting criteria, giving the weights directly is almost never feasible. Instead, a common method is to apply pairwise comparisons. Answers to the questions “How many times is Criterion A more important than Criterion B?” and so on (which are explicit cardinal ratios) can be arranged in a matrix, called a pairwise comparison matrix (PCM). Formally, a PCM is a square matrix $\mathbf{A} = [a_{ij}]_{i,j=1,\dots,n}$ with the properties $a_{ij} > 0$ and $a_{ij} = 1/a_{ji}$ (which implies $a_{ii} = 1$). If the cardinal transitivity property $a_{ik}a_{kj} = a_{ij} \ \forall i, j, k = 1, \dots, n$ also holds for a PCM, it is called consistent, otherwise it is called inconsistent [20]. Let $\mathcal{PCM}_n$ denote the set of PCMs of size $n \times n$. The next step is to extract the weights of criteria from the PCM. Several methods exist for this task [3, 9, 12, 17], but we will only focus on the most commonly used one, the eigenvector method (EM). The eigenvector method gives the weight vector $\mathbf{w}^{EM} = (w_1, \dots, w_n)^T$ as the right Perron eigenvector of $\mathbf{A} \in \mathcal{PCM}_n$, thus $\mathbf{A}\mathbf{w}^{EM} = \lambda_{\max}\mathbf{w}^{EM}$ holds, where $\lambda_{\max}$ is the principal eigenvalue of $\mathbf{A}$. $\lambda_{\max} \geq n$, and $\lambda_{\max} = n$ if and only if $\mathbf{A}$ is consistent [20]. A consistent PCM can be written as

$$\mathbf{A} = \begin{pmatrix} 1 & x_1 & x_2 & \dots & x_{n-1} \\ 1/x_1 & 1 & x_2/x_1 & \dots & x_{n-1}/x_1 \\ 1/x_2 & x_1/x_2 & 1 & \dots & x_{n-1}/x_2 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1/x_{n-1} & x_1/x_{n-1} & x_2/x_{n-1} & \dots & 1 \end{pmatrix} \in \mathcal{PCM}_n,$$

where $x_1, \dots, x_{n-1} > 0$.

The elements of a PCM approximate the ratios of the weights, therefore the ratios of the elements of the weight vector should be as close as possible to the corresponding matrix elements. If a weight vector cannot be trivially improved in this regard (there is no other weight vector which is at least as good approximation, and strictly better in at least one position), it is called Pareto optimal or efficient. It has been proved that the eigenvector method does not always produce an efficient solution. However, in some special cases the eigenvector method always gives an efficient weight vector. If the PCM is simple perturbed, i.e., it differs from a consistent PCM in only one element and its reciprocal, the principal right eigenvector is efficient [1]. In the paper this will be extended to double perturbed PCMs, which only differ from a consistent PCM in two elements and their reciprocals. These special types of PCMs are not just theoretically important, but also occur in real decision problems. Poesz [19] gathered a handful of empirical PCMs that were analyzed in [7]; out of 90 matrices of size at most 6×6 , 53 were either consistent, simple perturbed or double perturbed (see [7, Table 1]).

In Section 2 we will introduce the key definitions and tools used in the paper, together with an example. In Section 3 the main results of the paper will be proved: through obtaining explicit formulas for the principal right eigenvector and a series of lemmas, efficiency of the principal right eigenvector will be shown for the case of double perturbed PCMs. The proofs of the lemmas, except for one, are given in detail in the Appendix. In Section 4 conclusions follow.

2 Efficiency and perturbed pairwise comparison matrices

Pareto optimality or efficiency [13, Chapter 2][22, Chapter 6] is a basic concept of multi-objective optimization and multi-attribute decision making, too. The definition for weight vectors of PCM is as follows. Let $\mathbf{A} = [a_{ij}]_{i,j=1,\dots,n} \in \mathcal{PCM}_n$ and $\mathbf{w} = (w_1, w_2, \dots, w_n)^T$ be a positive weight vector.

Definition 1. A positive weight vector $\mathbf{w}$ is called *efficient* if no other positive weight vector $\mathbf{w}' = (w'_1, w'_2, \dots, w'_n)^T$ exists such that

$$\left| a_{ij} - \frac{w'_i}{w'_j} \right| \leq \left| a_{ij} - \frac{w_i}{w_j} \right| \quad \text{for all } 1 \leq i, j \leq n, \quad (1)$$

$$\left| a_{k\ell} - \frac{w'_k}{w'_\ell} \right| < \left| a_{k\ell} - \frac{w_k}{w_\ell} \right| \quad \text{for some } 1 \leq k, \ell \leq n. \quad (2)$$

A weight vector $\mathbf{w}$ is called *inefficient* if it is not efficient.

For a consistent PCM $a_{ij} = w_i^{EM}/w_j^{EM}$ [20], which implies the following remark:

Remark 1. The principal right eigenvector $\mathbf{w}^{EM}$ is efficient for every consistent PCM.

For inconsistent PCMs however, the principal right eigenvector can be inefficient, found by Blanquero, Carrizosa and Conde [4, Section 3]. This result was also reinforced by Bajwa, Choo and Wedley [3], by Conde and Pérez [10] and by Fedrizzi [16]. Blanquero, Carrizosa and Conde [4] developed LP models to test whether a weight vector is efficient. Bozóki and Fülöp [6] further developed the models and provided algorithms to improve an inefficient weight vector. Anholcer and Fülöp [2] devised a new algorithm to derive an efficient solution from an inconsistent PCM.

Furthermore, Bozóki [5] showed that the principal right eigenvector of a whole class of matrices, namely the parametric PCM

$$\mathbf{A}(p, q) = \begin{pmatrix} 1 & p & p & p & \dots & p & p \\ 1/p & 1 & q & 1 & \dots & 1 & 1/q \\ 1/p & 1/q & 1 & q & \dots & 1 & 1 \\ \vdots & \vdots & \vdots & \ddots & & \vdots & \vdots \\ \vdots & \vdots & \vdots & & \ddots & \vdots & \vdots \\ 1/p & 1 & 1 & 1 & \dots & 1 & q \\ 1/p & q & 1 & 1 & \dots & 1/q & 1 \end{pmatrix} \in \mathcal{PCM}_n,$$

where $n \geq 4$, $p > 0$ and $1 \neq q > 0$, is inefficient.

Several necessary and sufficient conditions were examined by Blanquero, Carrizosa and Conde [4], one of which is of crucial importance to us. This one uses a directed graph representation:

Definition 2. Let $\mathbf{A} = [a_{ij}]_{i,j=1,\dots,n} \in \mathcal{PCM}_n$ and $\mathbf{w} = (w_1, w_2, \dots, w_n)^T$ be a positive weight vector. A directed graph $G = (V, \vec{E})_{\mathbf{A}, \mathbf{w}}$ is defined as follows: $V = \{1, 2, \dots, n\}$ and

$$\vec{E} = \left\{ \text{arc}(i \rightarrow j) \left| \frac{w_i}{w_j} \geq a_{ij}, i \neq j \right. \right\}.$$

It follows from Definition 2 that if $w_i/w_j = a_{ij}$, then there is a bidirected arc between nodes i, j . The result of Blanquero, Carrizosa and Conde using this representation is as follows:

Theorem 1 ([4, Corollary 10]). *Let $\mathbf{A} \in \mathcal{PCM}_n$. A weight vector $\mathbf{w}$ is efficient if and only if $G = (V, \vec{E})_{\mathbf{A}, \mathbf{w}}$ is a strongly connected digraph, that is, there exist directed paths from i to j and from j to i for all pairs of nodes i, j .*

The following numerical example provides an illustration for Theorem 1.

Example 1. Let $\mathbf{A} \in \mathcal{PCM}_4$ be defined as follows:

$$\mathbf{A} = \begin{pmatrix} 1 & 1/2 & 4 & 2 \\ 2 & 1 & 5 & 7 \\ 1/4 & 1/5 & 1 & 2 \\ 1/2 & 1/7 & 1/2 & 1 \end{pmatrix}.$$

The principal right eigenvector $\mathbf{w}^{EM}$, corresponding to PCM $\mathbf{A}$, and the consistent approximation of $\mathbf{A}$ as generated by $\mathbf{w}^{EM}$ are as follows (truncated at 8 and 4 correct digits, respectively):

$$\mathbf{w}^{EM} = \begin{pmatrix} 0.27471631 \\ 0.53204485 \\ 0.10869376 \\ 0.08454506 \end{pmatrix}, \quad \left[\frac{w_i^{EM}}{w_j^{EM}} \right] = \begin{pmatrix} 1 & 0.5163 & 2.5274 & 3.2493 \\ 1.9367 & 1 & 4.8948 & 6.2930 \\ 0.3956 & 0.2042 & 1 & 1.2856 \\ 0.3077 & 0.1589 & 0.7778 & 1 \end{pmatrix}.$$

Apply Definition 2, the directed graph $G = (V, \vec{E})_{\mathbf{A}, \mathbf{w}^{EM}}$ corresponding to $\mathbf{A}$ and $\mathbf{w}^{EM}$ is drawn in Figure 1. By Theorem 1, $\mathbf{w}^{EM}$ is not efficient, because the corresponding digraph is not strongly connected: no arc leaves node 2.




Figure 1: The principal right eigenvector in Example 1 is inefficient, because the corresponding digraph is not strongly connected: no arc leaves node 2

It can be seen in a more direct way why the principal right eigenvector $\mathbf{w}^{EM}$ is inefficient. If the second coordinate is increased to $w'_2 \approx 0.54346883$ (truncated at 8 digits), then

$$\mathbf{w}' = \begin{pmatrix} 0.27471631 \\ 0.54346883 \\ 0.10869376 \\ 0.08454506 \end{pmatrix}, \quad \left[\frac{w'_i}{w'_j} \right] = \begin{pmatrix} 1 & \mathbf{0.5054} & 2.5274 & 3.2493 \\ \mathbf{1.9782} & 1 & \mathbf{5} & \mathbf{6.4281} \\ 0.3956 & \mathbf{0.2} & 1 & 1.2856 \\ 0.3077 & \mathbf{0.1555} & 0.7778 & 1 \end{pmatrix}.$$

The approximation in the entries marked by bold became strictly better ((2) holds in Definition 1), while for all other entries the approximation remained the same ((1) holds with equality in Definition 1).

As it can be seen from Example 1 above, Theorem 1 is a powerful and applicable characterization of efficiency.

In the rest of the paper, special types of PCM's are considered.

A *simple perturbed* PCM differs from a consistent PCM in only one element and its reciprocal, or in other words it can be made consistent by altering only one element (and its reciprocal). Thus, without loss of generality, a simple perturbed PCM can be written as

$$\mathbf{A}_\delta = \begin{pmatrix} 1 & \delta x_1 & x_2 & \dots & x_{n-1} \\ 1/(\delta x_1) & 1 & x_2/x_1 & \dots & x_{n-1}/x_1 \\ 1/x_2 & x_1/x_2 & 1 & \dots & x_{n-1}/x_2 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1/x_{n-1} & x_1/x_{n-1} & x_2/x_{n-1} & \dots & 1 \end{pmatrix} \in \mathcal{PCM}_n,$$

where $x_1, \dots, x_{n-1} > 0$ and $0 < \delta \neq 1$.

Theorem 2 ([1, Theorem 3.1]). *The principal right eigenvector of a simple perturbed pairwise comparison matrix is efficient.*

Similarly, a *double perturbed* PCM differs from a consistent PCM in two elements and their reciprocals, or in other words it can be made consistent by altering two elements (and their reciprocals). We have to differentiate between three cases of double perturbed PCM's. Without loss of generality, every double perturbed PCM is equivalent to one of them. Also, we can suppose without the loss of generality, that from now on $n \geq 4$, because a PCM with $n = 3$ is either simple perturbed or consistent. In Case 1, the perturbed elements are in the same row, and they are multiplied by $0 < \delta \neq 1$ and $0 < \gamma \neq 1$ respectively. In Case 2, they are in different rows, but this case needs to be further divided into two subcases (2A and 2B) due to algebraic issues. In Case 2A matrix size is 4×4 , while in Case 2B matrix size is at least 5×5 . Thus, these matrices take the following form:

Case 1:

$$\mathbf{P}_{\gamma, \delta} = \begin{pmatrix} 1 & \delta x_1 & \gamma x_2 & x_3 & \dots & x_{n-1} \\ 1/(\delta x_1) & 1 & x_2/x_1 & x_3/x_1 & \dots & x_{n-1}/x_1 \\ 1/(\gamma x_2) & x_1/x_2 & 1 & x_3/x_2 & \dots & x_{n-1}/x_2 \\ 1/x_3 & x_1/x_3 & x_2/x_3 & 1 & \dots & x_{n-1}/x_3 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 1/x_{n-1} & x_1/x_{n-1} & x_2/x_{n-1} & x_3/x_{n-1} & \dots & 1 \end{pmatrix}, \quad (3)$$

Case 2A:

$$\mathbf{Q}_{\gamma, \delta} = \begin{pmatrix} 1 & \delta x_1 & x_2 & x_3 \\ 1/(\delta x_1) & 1 & x_2/x_1 & x_3/x_1 \\ 1/x_2 & x_1/x_2 & 1 & \gamma x_3/x_2 \\ 1/x_3 & x_1/x_3 & x_2/(\gamma x_3) & 1 \end{pmatrix}, \quad (4)$$

Case 2B:

$$\mathbf{R}_{\gamma,\delta} = \begin{pmatrix} 1 & \delta x_1 & x_2 & x_3 & x_4 & \dots & x_{n-1} \\ 1/(\delta x_1) & 1 & x_2/x_1 & x_3/x_1 & x_4/x_1 & \dots & x_{n-1}/x_1 \\ 1/x_2 & x_1/x_2 & 1 & \gamma x_3/x_2 & x_4/x_2 & \dots & x_{n-1}/x_2 \\ 1/x_3 & x_1/x_3 & x_2/(\gamma x_3) & 1 & x_4/x_3 & \dots & x_{n-1}/x_3 \\ 1/x_4 & x_1/x_4 & x_2/x_4 & x_3/x_4 & 1 & \dots & x_{n-1}/x_4 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 1/x_{n-1} & x_1/x_{n-1} & x_2/x_{n-1} & x_3/x_{n-1} & x_4/x_{n-1} & \dots & 1 \end{pmatrix}. \quad (5)$$

Once again, $x_1, \dots, x_{n-1} > 0$ and $0 < \delta, \gamma \neq 1$.

Remark 2. If either $\delta = 1$ or $\gamma = 1$ then the PCM is simple perturbed. If $\delta = \gamma = 1$ then the PCM is consistent.

Remark 3. If $n = 4$ and $\delta = \gamma$, then the PCM $\mathbf{P}_{\delta,\delta}$ in Case 1 is simple perturbed (multiply the single element x_3 in position (1,4) by δ to have a consistent PCM).

Bozóki, Fülöp and Poesz examined PCMs that can be made consistent by modifying at most 3 elements [7]. Each of the three cases above corresponds to a graph: Case 1 corresponds to [7, Fig. 6(b)] while Case 2 corresponds to [7, Fig. 6(a)]. Cook and Kress [11] and Brunelli and Fedrizzi [8] also examined the similar idea of comparing two PCMs that differ in only one element.

3 Main results

The main result of the paper will be the extension of Theorem 2 for double perturbed PCMs. A method to acquire the explicit form of the principal right eigenvector of a PCM when the perturbed elements are in the same row or column has been developed by Farkas, Rózsa and Stubnya [15]. Farkas [14] writes the explicit formula for the simple perturbed case. Our first goal is to extend the method for the double perturbed case. Similar to [14], the characteristic polynomial is needed first.

Let $\mathbf{D} = \text{diag}(1, 1/x_1, \dots, 1/x_{n-1})$, and let $\mathbf{e} = (1, \dots, 1)^T$. For $\mathbf{A} \in \mathcal{PCM}_n$,

$$\mathbf{e}\mathbf{e}^T - \mathbf{U}_i \mathbf{V}_i^T = \mathbf{D}^{-1} \mathbf{A} \mathbf{D} \quad (6)$$

holds for $i = 1, 2$, where

$$\mathbf{U}_1 = \begin{pmatrix} 0 & 1 \\ 1 - 1/\delta & 0 \\ 1 - 1/\gamma & 0 \\ 0 & 0 \\ \vdots & \vdots \\ 0 & 0 \end{pmatrix} \in \mathbb{R}^{n \times 2}, \quad \mathbf{V}_1 = \begin{pmatrix} 1 & 0 \\ 0 & 1 - \delta \\ 0 & 1 - \gamma \\ 0 & 0 \\ \vdots & \vdots \\ 0 & 0 \end{pmatrix} \in \mathbb{R}^{n \times 2}, \quad (7)$$

$$\mathbf{U}_2 = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 - 1/\delta & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 - 1/\gamma & 0 \\ 0 & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 \end{pmatrix} \in \mathbb{R}^{n \times 4}, \quad \mathbf{V}_2 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 - \delta & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 - \gamma \\ 0 & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 \end{pmatrix} \in \mathbb{R}^{n \times 4}. \quad (8)$$

Case 1 with form (3) corresponds to $i = 1$, while Case 2 with forms (4) and (5) corresponds to $i = 2$.

Lemma 1 (Matrix determinant lemma, [18]). *If $\mathbf{A} \in \mathbb{R}^{n \times n}$ is invertible, and $\mathbf{U}, \mathbf{V} \in \mathbb{R}^{n \times m}$, then*

$$\det(\mathbf{A} + \mathbf{U}\mathbf{V}^T) = \det(\mathbf{I}_m + \mathbf{V}^T \mathbf{A}^{-1} \mathbf{U}) \det(\mathbf{A}),$$

where $\mathbf{I}_m$ denotes the identity matrix of size $m \times m$.

Lemma 2 (Sherman–Morrison formula, [21]). *Let $\mathbf{A} \in \mathbb{R}^{n \times n}$, $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$. If $\mathbf{A}$ is invertible and $1 + \mathbf{v}^T \mathbf{A}^{-1} \mathbf{u} \neq 0$, then $(\mathbf{A} + \mathbf{u}\mathbf{v}^T)^{-1}$ exists, and*

$$(\mathbf{A} + \mathbf{u}\mathbf{v}^T)^{-1} = \mathbf{A}^{-1} - \frac{1}{1 + \mathbf{v}^T \mathbf{A}^{-1} \mathbf{u}} \mathbf{A}^{-1} \mathbf{u} \mathbf{v}^T \mathbf{A}^{-1}.$$

Let $\mathbf{A} \in \mathcal{PCM}_n$ be a double perturbed PCM and $\mathbf{U}_i, \mathbf{V}_i$ be as in (6). Let the matrix $\mathbf{K}_{\mathbf{A}}(\lambda) \in \mathbb{R}^{n \times n}$ be defined as follows:

$$\mathbf{K}_{\mathbf{A}}(\lambda) = \lambda \mathbf{I} + \mathbf{U}_i \mathbf{V}_i^T - \mathbf{e} \mathbf{e}^T = \lambda \mathbf{I} - \mathbf{D}^{-1} \mathbf{A} \mathbf{D},$$

where $\mathbf{I}$ denotes $\mathbf{I}_n$, $\mathbf{e} = (1, \dots, 1)^T \in \mathbb{R}^n$, $i = 1$ in Case 1, $i = 2$ in Case 2, and the second equation follows from (6).

Lemma 3. *The characteristic polynomial of the double perturbed PCM $\mathbf{A} \in \mathcal{PCM}_n$ is*

$$p_{\mathbf{A}}(\lambda) = (-1)^n \det(\mathbf{K}_{\mathbf{A}}(\lambda)).$$

Proof. As before, $i = 1$ in Case 1 and $i = 2$ in Case 2.

$$\begin{aligned} p_{\mathbf{A}}(\lambda) &= \det(\mathbf{A} - \lambda \mathbf{I}) \\ &= (-1)^n \det(\lambda \mathbf{I} - \mathbf{A}) \\ &= (-1)^n \det(\lambda \mathbf{I} + \mathbf{D} (\mathbf{U}_i \mathbf{V}_i^T - \mathbf{e} \mathbf{e}^T) \mathbf{D}^{-1}) \\ &= (-1)^n \det(\mathbf{D} (\lambda \mathbf{I} + \mathbf{U}_i \mathbf{V}_i^T - \mathbf{e} \mathbf{e}^T) \mathbf{D}^{-1}) \\ &= (-1)^n \det(\mathbf{D}) \det(\lambda \mathbf{I} + \mathbf{U}_i \mathbf{V}_i^T - \mathbf{e} \mathbf{e}^T) \det(\mathbf{D}^{-1}) \\ &= (-1)^n \det(\mathbf{K}_{\mathbf{A}}(\lambda)). \end{aligned}$$

□

Lemma 4. $\det(\lambda \mathbf{I} - \mathbf{e} \mathbf{e}^T) = \lambda^n - n \lambda^{n-1}$.

Proof. If $\lambda = 0$, then both sides of the equation are 0. If $\lambda \neq 0$, apply Lemma 1 with $m = 1$, $\mathbf{A} = \lambda \mathbf{I}$, $\mathbf{U} = -\mathbf{e}$, $\mathbf{V} = \mathbf{e}$:

$$\det(\lambda \mathbf{I} - \mathbf{e} \mathbf{e}^T) = (1 - \mathbf{e}^T (\lambda \mathbf{I})^{-1} \mathbf{e}) \det(\lambda \mathbf{I}) = \lambda^n - n \lambda^{n-1}.$$

□

Lemma 5. *If $\lambda \neq 0$ and $\lambda \neq n$, then $(\lambda \mathbf{I} - \mathbf{e} \mathbf{e}^T)^{-1}$ exists, and*

$$(\lambda \mathbf{I} - \mathbf{e} \mathbf{e}^T)^{-1} = \frac{1}{\lambda(\lambda - n)} \mathbf{e} \mathbf{e}^T + \frac{1}{\lambda} \mathbf{I}.$$

Proof. Apply the Sherman–Morrison formula (Lemma 2) with $\mathbf{A} = \lambda \mathbf{I}$, $\mathbf{u} = -\mathbf{e}$, $\mathbf{v} = \mathbf{e}$. $\square$

Lemma 6. Let $\mathbf{U}, \mathbf{V} \in \mathbb{R}^{n \times m}$ be arbitrary matrices. If $\lambda \neq 0$ and $\lambda \neq n$, then

$$\det(\lambda \mathbf{I}_n + \mathbf{U} \mathbf{V}^T - \mathbf{e} \mathbf{e}^T) = (\lambda^n - n \lambda^{n-1}) \det\left(\mathbf{I}_m + \frac{1}{\lambda(\lambda - n)} \mathbf{V}^T \mathbf{e} \mathbf{e}^T \mathbf{U} + \frac{1}{\lambda} \mathbf{V}^T \mathbf{U}\right).$$

Proof. Apply Lemma 1 with $\mathbf{A} = \lambda \mathbf{I}_n - \mathbf{e} \mathbf{e}^T$. According to Lemma 5, $\mathbf{A}$ is invertible. Utilizing Lemmas 1, 4 and 5 the following equations hold:

$$\begin{aligned} & \det((\lambda \mathbf{I}_n - \mathbf{e} \mathbf{e}^T) + \mathbf{U} \mathbf{V}^T) \\ &= \det\left(\mathbf{I}_m + \mathbf{V}^T (\lambda \mathbf{I}_n - \mathbf{e} \mathbf{e}^T)^{-1} \mathbf{U}\right) \det(\lambda \mathbf{I}_n - \mathbf{e} \mathbf{e}^T) \\ &= \det\left(\mathbf{I}_m + \frac{1}{\lambda(\lambda - n)} \mathbf{V}^T \mathbf{e} \mathbf{e}^T \mathbf{U} + \frac{1}{\lambda} \mathbf{V}^T \mathbf{U}\right) (\lambda^n - n \lambda^{n-1}). \end{aligned}$$

$\square$

We can write the characteristic polynomial of double perturbed PCM in explicit form.

Theorem 3. Let $n \geq 4$. The characteristic polynomial of a double perturbed PCM in form (3) (Case 1) is

$$p_{\mathbf{P}}(\lambda) = (-1)^n \lambda^{n-3} \left(\lambda^3 - n \lambda^2 - \left(\frac{\gamma}{\delta} + \frac{\delta}{\gamma} \right) - (n-3) \left(\gamma + \delta + \frac{1}{\gamma} + \frac{1}{\delta} \right) + 4n - 10 \right).$$

Proof. Lemma 3 implies that

$$p_{\mathbf{P}}(\lambda) = (-1)^n \det(\mathbf{K}_{\mathbf{P}}(\lambda)) = (-1)^n \det(\lambda \mathbf{I} + \mathbf{U}_1 \mathbf{V}_1^T - \mathbf{e} \mathbf{e}^T),$$

where $\mathbf{U}_1$ and $\mathbf{V}_1$ are defined by (7). Suppose that $\lambda \neq n$ and $\lambda \neq 0$. According to Lemma 6

$$\begin{aligned} p_{\mathbf{P}}(\lambda) &= (-1)^n (\lambda^n - n \lambda^{n-1}) \det\left(\mathbf{I}_2 + \frac{1}{\lambda(\lambda - n)} \mathbf{V}_1^T \mathbf{e} \mathbf{e}^T \mathbf{U}_1 + \frac{1}{\lambda} \mathbf{V}_1^T \mathbf{U}_1\right) \\ &= (-1)^n (\lambda^n - n \lambda^{n-1}) \det(\mathbf{S}) \\ &= (-1)^n \lambda^{n-3} \left(\lambda^3 - n \lambda^2 - \left(\frac{\gamma}{\delta} + \frac{\delta}{\gamma} \right) - (n-3) \left(\gamma + \delta + \frac{1}{\gamma} + \frac{1}{\delta} \right) + 4n - 10 \right), \end{aligned}$$

where

$$\mathbf{S} = \begin{pmatrix} 1 + \frac{2-1/\delta-1/\gamma}{\lambda(\lambda-n)} & \frac{1}{\lambda(\lambda-n)} + \frac{1}{\lambda} \\ \frac{(2-\delta-\gamma)(1-1/\delta)+(2-\delta-\gamma)(1-1/\gamma)}{\lambda(\lambda-n)} + \frac{(1-\delta)(1-1/\delta)}{\lambda} + \frac{(1-\gamma)(1-1/\gamma)}{\lambda} & 1 + \frac{2-\delta-\gamma}{\lambda(\lambda-n)} \end{pmatrix}.$$

A polynomial of degree n is uniquely determined by $n+1$ points, and we have calculated $p_{\mathbf{P}}(\lambda)$ in all but two points, which completes the proof. $\square$

Theorem 4. Let $n \geq 4$. The characteristic polynomial of a double perturbed PCM in form (5) (Case 2B) is

$$p_{\mathbf{R}}(\lambda) = (-1)^n \lambda^{n-5} \left(\lambda^5 - n\lambda^4 - (n-2) \left(\gamma + \delta + \frac{1}{\gamma} + \frac{1}{\delta} - 4 \right) \lambda^2 - c\lambda - (n-4)c \right),$$

where

$$c = \frac{(\gamma-1)^2(\delta-1)^2}{\gamma\delta}.$$

Furthermore, the characteristic polynomial of a double perturbed PCM in form (4) (Case 2A), $p_{\mathbf{Q}}(\lambda)$ is a special case of $p_{\mathbf{R}}(\lambda)$ with $n = 4$. Namely,

$$p_{\mathbf{Q}}(\lambda) = \lambda^4 - 4\lambda^3 - 2 \left(\gamma + \delta + \frac{1}{\gamma} + \frac{1}{\delta} - 4 \right) \lambda - \frac{(\gamma-1)^2(\delta-1)^2}{\gamma\delta}.$$

Proof. Lemma 3 implies that

$$p_{\mathbf{R}}(\lambda) = (-1)^n \det(\mathbf{K}_{\mathbf{R}}(\lambda)) = (-1)^n \det(\lambda \mathbf{I} + \mathbf{U}_2 \mathbf{V}_2^T - \mathbf{e} \mathbf{e}^T),$$

where $\mathbf{U}_2$ and $\mathbf{V}_2$ are defined by (8). Suppose that $\lambda \neq n$ and $\lambda \neq 0$. According to Lemma 6

$$\begin{aligned} p_{\mathbf{R}}(\lambda) &= (-1)^n (\lambda^n - n\lambda^{n-1}) \det \left(\mathbf{I}_4 + \frac{1}{\lambda(\lambda-n)} \mathbf{V}_2^T \mathbf{e} \mathbf{e}^T \mathbf{U}_2 + \frac{1}{\lambda} \mathbf{V}_2^T \mathbf{U}_2 \right) \\ &= (-1)^n (\lambda^n - n\lambda^{n-1}) \det(\mathbf{T}) \\ &= (-1)^n \lambda^{n-5} \left(\lambda^5 - n\lambda^4 - (n-2) \left(\gamma + \delta + \frac{1}{\gamma} + \frac{1}{\delta} - 4 \right) \lambda^2 - c\lambda - (n-4)c \right), \end{aligned}$$

where

$$\mathbf{T} = \begin{pmatrix} 1 + \frac{1-1/\delta}{\lambda(\lambda-n)} & \frac{1}{\lambda(\lambda-n)} + \frac{1}{\lambda} & \frac{1-1/\gamma}{\lambda(\lambda-n)} & \frac{1}{\lambda(\lambda-n)} \\ \frac{(1-\delta)(1-1/\delta)}{\lambda(\lambda-n)} + \frac{(1-\delta)(1-1/\delta)}{\lambda} & 1 + \frac{1-\delta}{\lambda(\lambda-n)} & \frac{(1-\delta)(1-1/\gamma)}{\lambda(\lambda-n)} & \frac{1-\delta}{\lambda(\lambda-n)} \\ \frac{1-1/\delta}{\lambda(\lambda-n)} & \frac{1}{\lambda(\lambda-n)} & 1 + \frac{1-1/\gamma}{\lambda(\lambda-n)} & \frac{1}{\lambda(\lambda-n)} + \frac{1}{\lambda} \\ \frac{(1-\gamma)(1-1/\delta)}{\lambda(\lambda-n)} & \frac{1-\gamma}{\lambda(\lambda-n)} & \frac{(1-\gamma)(1-1/\gamma)}{\lambda(\lambda-n)} + \frac{(1-\gamma)(1-1/\gamma)}{\lambda} & 1 + \frac{1-\gamma}{\lambda(\lambda-n)} \end{pmatrix}$$

and

$$c = \frac{(\gamma-1)^2(\delta-1)^2}{\gamma\delta}.$$

Again, a polynomial of degree n is uniquely determined by $n+1$ points, and we have calculated $p_{\mathbf{R}}(\lambda)$ in all but two points, which completes the proof. The case $n = 4$ is analogous, and

$$p_{\mathbf{Q}}(\lambda) = \lambda^4 - 4\lambda^3 - 2 \left(\gamma + \delta + \frac{1}{\gamma} + \frac{1}{\delta} - 4 \right) \lambda - \frac{(\gamma-1)^2(\delta-1)^2}{\gamma\delta}$$

is resulted in. □

Theorem 5. *The principle right eigenvector of a double perturbed PCM can be written in explicit ways.*

In Case 1 (γ and δ are in the same row), the formulas for the principal right eigenvector are the following:

$$\mathbf{w}^{EM} = \begin{pmatrix} \delta\gamma\lambda(\lambda - n + 1) \\ \frac{1}{x_1} [\gamma\lambda - (n-2)\gamma + \delta + (n-3)\delta\gamma] \\ \frac{1}{x_2} [\delta\lambda - (n-2)\delta + \gamma + (n-3)\delta\gamma] \\ \frac{1}{x_3} [\gamma + \delta + \delta\gamma\lambda - 2\delta\gamma] \\ \vdots \\ \frac{1}{x_{i-1}} [\gamma + \delta + \delta\gamma\lambda - 2\delta\gamma] \\ \vdots \\ \frac{1}{x_{n-1}} [\gamma + \delta + \delta\gamma\lambda - 2\delta\gamma] \end{pmatrix}, \quad (9)$$

$$\mathbf{w}^{EM} = c_{11} \begin{pmatrix} x_1\gamma\lambda [\delta\lambda - (n-2)\delta + \gamma + n-3] \\ \gamma\lambda^3 - (n-1)\gamma\lambda^2 - (n-3)(\gamma^2 - 2\gamma + 1) \\ \frac{x_1}{x_2} [\gamma\lambda^2 - \gamma\lambda + \delta\lambda + (n-3)(\delta\gamma - \delta - \gamma + 1)] \\ \frac{x_1}{x_3} [\gamma\lambda^2 - \gamma\lambda - \gamma + \delta + \delta\gamma\lambda - \delta\gamma + \gamma^2] \\ \vdots \\ \frac{x_1}{x_{i-1}} [\gamma\lambda^2 - \gamma\lambda - \gamma + \delta + \delta\gamma\lambda - \delta\gamma + \gamma^2] \\ \vdots \\ \frac{x_1}{x_{n-1}} [\gamma\lambda^2 - \gamma\lambda - \gamma + \delta + \delta\gamma\lambda - \delta\gamma + \gamma^2] \end{pmatrix}, \quad (10)$$

$$\mathbf{w}^{EM} = c_{12} \begin{pmatrix} x_2\delta\lambda [\delta + \gamma\lambda - (n-2)\gamma + n-3] \\ \frac{x_2}{x_1} [\delta\lambda^2 - \delta\lambda + \gamma\lambda + (n-3)(\delta\gamma - \delta - \gamma + 1)] \\ \delta\lambda^3 - (n-1)\delta\lambda^2 - (n-3)(\delta^2 - 2\delta + 1) \\ \frac{x_2}{x_3} [\delta\lambda^2 - \delta\lambda + \gamma - \delta + \delta^2 + \delta\gamma\lambda - \delta\gamma] \\ \vdots \\ \frac{x_2}{x_{i-1}} [\delta\lambda^2 - \delta\lambda + \gamma - \delta + \delta^2 + \delta\gamma\lambda - \delta\gamma] \\ \vdots \\ \frac{x_2}{x_{n-1}} [\delta\lambda^2 - \delta\lambda + \gamma - \delta + \delta^2 + \delta\gamma\lambda - \delta\gamma] \end{pmatrix}, \quad (11)$$

$$\mathbf{w}^{EM} = c_{13} \begin{pmatrix} x_3\delta\gamma\lambda(\delta + \gamma + \lambda - 2) \\ \frac{x_3}{x_1} [\delta\gamma\lambda^2 - \delta\gamma\lambda + \gamma^2 + \gamma\lambda - \gamma - \delta\gamma + \delta] \\ \frac{x_3}{x_2} [\delta\gamma\lambda^2 - \delta\gamma\lambda - \delta\gamma + \gamma + \delta^2 + \delta\lambda - \delta] \\ \delta\gamma\lambda^2 - 4\delta\gamma + \gamma + \delta + \delta^2\gamma + \gamma^2\delta \\ \frac{x_3}{x_4} [\delta\gamma\lambda^2 - 4\delta\gamma + \gamma + \delta + \delta^2\gamma + \gamma^2\delta] \\ \vdots \\ \frac{x_3}{x_{n-1}} [\delta\gamma\lambda^2 - 4\delta\gamma + \gamma + \delta + \delta^2\gamma + \gamma^2\delta] \end{pmatrix}, \quad (12)$$

where $c_{11}, c_{12}, c_{13} \in \mathbb{R}$.

In Case 2A (γ and δ are in different rows, and matrix size is 4×4) the formulas take the following form:

$$\mathbf{w}^{EM} = \begin{pmatrix} \delta(\lambda^3\gamma - 3\lambda^2\gamma - 1 + 2\gamma - \gamma^2) \\ \frac{1}{x_1} [\lambda^2\gamma - 2\lambda\gamma + \delta + 2\lambda\delta\gamma - 2\delta\gamma + \delta\gamma^2] \\ \frac{1}{x_2} \gamma [\gamma + \lambda - 1 + \delta\lambda^2 - 2\lambda\delta + \delta + \lambda\delta\gamma - \delta\gamma] \\ \frac{1}{x_3} [1 + \lambda\gamma - \gamma + \lambda\delta - \delta + \delta\gamma\lambda^2 - 2\lambda\delta\gamma + \delta\gamma] \end{pmatrix}, \quad (13)$$

$$\mathbf{w}^{EM} = c_{21} \begin{pmatrix} x_1[\delta\gamma\lambda^2 - 2\lambda\delta\gamma + 1 + 2\lambda\gamma - 2\gamma + \gamma^2] \\ \lambda^3\gamma - 3\lambda^2\gamma - 1 + 2\gamma - \gamma^2 \\ \frac{x_1}{x_2} \gamma [\lambda\gamma + \lambda^2 - 2\lambda - \gamma + 1 + \lambda\delta - \delta + \delta\gamma] \\ \frac{x_1}{x_3} [\lambda + \lambda^2\gamma - 2\lambda\gamma - 1 + \gamma + \delta + \lambda\delta\gamma - \delta\gamma] \end{pmatrix}, \quad (14)$$

$$\mathbf{w}^{EM} = c_{22} \begin{pmatrix} x_2\delta(1 + \lambda\gamma - \gamma)(\delta + \lambda - 1) \\ \frac{x_2}{x_1} [1 + \lambda\gamma - \gamma + \lambda\delta - \delta + \delta\gamma\lambda^2 - 2\lambda\delta\gamma + \delta\gamma] \\ \gamma(\delta\lambda^3 - 3\delta\lambda^2 - 1 + 2\delta - \delta^2) \\ \frac{x_2}{x_3} [2\lambda\delta\gamma + \delta\lambda^2 - 2\lambda\delta - 2\delta\gamma + \gamma + \delta^2\gamma] \end{pmatrix}, \quad (15)$$

$$\mathbf{w}^{EM} = c_{23} \begin{pmatrix} x_3\delta(\lambda\gamma + \lambda^2 - 2\lambda - \gamma + 1 + \lambda\delta - \delta + \delta\gamma) \\ \frac{x_3}{x_1} [\gamma + \lambda - 1 + \delta\lambda^2 - 2\lambda\delta + \delta + \lambda\delta\gamma - \delta\gamma] \\ \frac{x_3}{x_2} [2\lambda\delta + \delta\gamma\lambda^2 - 2\lambda\delta\gamma - 2\delta + 1 + \delta^2] \\ \delta\lambda^3 - 3\delta\lambda^2 - 1 + 2\delta - \delta^2 \end{pmatrix}, \quad (16)$$

where $c_{21}, c_{22}, c_{23} \in \mathbb{R}$.

In Case 2B (γ and δ are in different rows, and matrix size is at least 5×5) the formulas are the following:

$$\mathbf{w}^{EM} = \begin{pmatrix} \delta\lambda[\lambda^3\gamma - (n-1)\lambda^2\gamma - (n-3)(\gamma^2 - 2\gamma + 1)] \\ \frac{1}{x_1} \{ \lambda^3\gamma - (n-2)\lambda^2\gamma + (n-2)\delta\gamma\lambda^2 + [\lambda\delta + (n-4)(\delta-1)](\gamma^2 - 2\gamma + 1) \} \\ \frac{1}{x_2} \gamma\lambda [\gamma + \lambda - 1 + \delta\lambda^2 - 2\lambda\delta + \delta + \lambda\delta\gamma - \delta\gamma] \\ \frac{1}{x_3} \lambda [1 + \lambda\gamma - \gamma + \lambda\delta - \delta + \delta\gamma\lambda^2 - 2\lambda\delta\gamma + \delta\gamma] \\ \frac{1}{x_4} [\gamma^2 - 2\gamma + \lambda^2\gamma + 1 + \lambda\delta - \delta\gamma\lambda^2 - 2\lambda\delta\gamma + \lambda\gamma^2\delta + \lambda^3\delta\gamma - \delta + 2\delta\gamma - \delta\gamma^2] \\ \vdots \\ \frac{1}{x_{n-1}} [\gamma^2 - 2\gamma + \lambda^2\gamma + 1 + \lambda\delta - \delta\gamma\lambda^2 - 2\lambda\delta\gamma + \lambda\gamma^2\delta + \lambda^3\delta\gamma - \delta + 2\delta\gamma - \delta\gamma^2] \end{pmatrix}, \quad (17)$$

$$\mathbf{w}^{EM} = c_{31} \begin{pmatrix} x_1[\lambda^3\delta\gamma - (n-2)\delta\gamma\lambda^2 - (n-4)\delta(\gamma-1)^2 + \lambda + (n-2)\lambda^2\gamma - 2\lambda\gamma + \lambda\gamma^2 + (n-4)(\gamma-1)^2] \\ \lambda(\lambda^3\gamma - (n-1)\lambda^2\gamma - (n-3)(\gamma-1)^2) \\ \frac{x_1}{x_2} \gamma\lambda(\lambda\gamma + \lambda^2 - 2\lambda - \gamma + 1 + \delta\lambda - \delta + \delta\gamma) \\ \frac{x_1}{x_3} \lambda(\lambda + \lambda^2\gamma - 2\lambda\gamma - 1 + \gamma + \delta + \lambda\delta\gamma - \delta\gamma) \\ \frac{x_1}{x_4} (\lambda\gamma^2 - 2\lambda\gamma + \lambda^3\gamma + \lambda - \gamma^2 + 2\gamma - \lambda^2\gamma - 1 + \delta - 2\delta\gamma + \delta\gamma^2 + \delta\gamma\lambda^2) \\ \vdots \\ \frac{x_1}{x_{n-1}} (\lambda\gamma^2 - 2\lambda\gamma + \lambda^3\gamma + \lambda - \gamma^2 + 2\gamma - \lambda^2\gamma - 1 + \delta - 2\delta\gamma + \delta\gamma^2 + \delta\gamma\lambda^2) \end{pmatrix}, \quad (18)$$

$$\mathbf{w}^{EM} = c_{32} \begin{pmatrix} x_2\delta\lambda(1 + \lambda\gamma - \gamma)(\delta + \lambda - 1) \\ \frac{x_2}{x_1} \lambda(1 + \lambda\gamma - \gamma)(1 + \delta\lambda - \delta) \\ \gamma\lambda[\lambda^3\delta - (n-1)\delta\lambda^2 - (n-3)(\delta-1)^2] \\ \frac{x_2}{x_3} [\delta\lambda^3 - (n-2)\delta\lambda^2(1-\gamma) - 2\lambda\delta\gamma + 2(n-4)\delta(1-\gamma) + \lambda\gamma + \delta^2\lambda\gamma + (n-4)(-1+\gamma-\delta^2+\delta^2\gamma)] \\ \frac{x_2}{x_4} (1 + \lambda\gamma - \gamma)(\delta\lambda^2 + 1 - 2\delta + \delta^2) \\ \vdots \\ \frac{x_2}{x_{n-1}} (1 + \lambda\gamma - \gamma)(\delta\lambda^2 + 1 - 2\delta + \delta^2) \end{pmatrix}, \quad (19)$$

$$\mathbf{w}^{EM} = c_{33} \begin{pmatrix} x_3 \delta \lambda (\lambda \gamma + \lambda^2 - 2\lambda - \gamma + 1 + \delta \lambda - \delta + \delta \gamma) \\ \frac{x_3}{x_1} \lambda (\gamma + \lambda - 1) (1 + \delta \lambda - \delta) \\ \frac{x_3}{x_2} [\lambda^3 \delta \gamma - (n-2) \delta \lambda^2 (\gamma - 1) - 2 \delta \lambda + 2(n-4) \delta (\gamma - 1) + \lambda + \delta^2 \lambda + (n-4) (1 - \gamma + \delta^2 - \delta^2 \gamma)] \\ \lambda [\delta \lambda^3 - (n-1) \delta \lambda^2 - (n-3) (\delta - 1)^2] \\ \frac{x_3}{x_4} (\delta \gamma \lambda^2 + \lambda^3 \delta - \delta \lambda^2 - 2 \delta \lambda - 2 \delta \gamma + 2 \delta - 1 + \gamma + \lambda + \delta^2 \lambda - \delta^2 + \delta^2 \gamma) \\ \vdots \\ \frac{x_3}{x_{n-1}} (\delta \gamma \lambda^2 + \lambda^3 \delta - \delta \lambda^2 - 2 \delta \lambda - 2 \delta \gamma + 2 \delta - 1 + \gamma + \lambda + \delta^2 \lambda - \delta^2 + \delta^2 \gamma) \end{pmatrix}, \quad (20)$$

$$\mathbf{w}^{EM} = c_{34} \begin{pmatrix} x_4 \delta \lambda (\gamma^2 - 2\gamma + \lambda^2 \gamma + 1) (\delta + \lambda - 1) \\ \frac{x_4}{x_1} \lambda (\gamma^2 - 2\gamma + \lambda^2 \gamma + 1) (1 + \delta \lambda - \delta) \\ \frac{x_4}{x_2} \gamma \lambda (\delta \gamma \lambda^2 + \lambda^3 \delta - \delta \lambda^2 - 2 \delta \lambda - 2 \delta \gamma + 2 \delta - 1 + \gamma + \lambda + \delta^2 \lambda - \delta^2 + \delta^2 \gamma) \\ \frac{x_4}{x_3} \lambda (\delta \lambda^2 + \lambda^3 \delta \gamma - \delta \gamma \lambda^2 - 2 \lambda \delta \gamma - 2 \delta + 2 \delta \gamma - \gamma + 1 + \lambda \gamma + \delta^2 + \delta^2 \lambda \gamma - \delta^2 \gamma) \\ (\gamma^2 - 2\gamma + \lambda^2 \gamma + 1) (\delta \lambda^2 + 1 - 2\delta + \delta^2) \\ \frac{x_4}{x_5} (\gamma^2 - 2\gamma + \lambda^2 \gamma + 1) (\delta \lambda^2 + 1 - 2\delta + \delta^2) \\ \vdots \\ \frac{x_4}{x_{n-1}} (\gamma^2 - 2\gamma + \lambda^2 \gamma + 1) (\delta \lambda^2 + 1 - 2\delta + \delta^2) \end{pmatrix}, \quad (21)$$

where $c_{31}, c_{32}, c_{33}, c_{34} \in \mathbb{R}$.

Proof. The proof is similar to that of the eigenvector formulas (24)–(26) in [14]. Let us consider Case 1. Let $\mathbf{D} = \text{diag}(1, 1/x_1, \dots, 1/x_{n-1})$, and let $\mathbf{K}_P(\lambda) = \lambda \mathbf{I} + \mathbf{U}_1 \mathbf{V}_1^T - \mathbf{e} \mathbf{e}^T$, with $\mathbf{U}_1$ and $\mathbf{V}_1$ as defined by (7). Since $\mathbf{D}$ is invertible, every column of the one rank matrix $\mathbf{D} \text{adj}(\mathbf{K}_P(\lambda_{\max})) \mathbf{D}^{-1}$ is a Perron eigenvector of $\mathbf{P}$. For Case 2, replace $\mathbf{U}_1$ by $\mathbf{U}_2$ and $\mathbf{V}_1$ by $\mathbf{V}_2$ as defined by (8). $\square$

The formulas for the Perron eigenvector have been written in different forms in Theorem 5. Although it is not at all apparent, these forms are equal up to a constant multiplier.

Using these formulas, the paper's main result can be obtained through a series of lemmas. Each of these lemmas corresponds to a directed edge in a digraph. Using these results, the direction of certain arcs can be determined. Thus, it will be shown that directed graphs of Cases 1, 2A and 2B are strongly connected. By Theorem 1, efficiency of the principal right eigenvector is implied.

It follows from the positivity of $\mathbf{w}^{EM}$ (see Remark 4 in the Appendix), that both sides of the starting inequalities of each lemma can be multiplied by the respective w_i^{EM} without further discussion. Since there are 28 lemmas, only one (Lemma 3f) has its proof in the main text as an illustrative example. The rest of the proofs are in the Appendix.

The first group of lemmas correspond to Case 1 (γ and δ are in the same row), i.e., the double perturbed PCM is written in form (3).

Lemma 1a. $\delta > 1$ and $\delta \geq \gamma \Rightarrow w_1^{EM}/w_2^{EM} < \delta x_1$.

Lemma 1b. $\delta < 1$ and $\delta \leq \gamma \Rightarrow w_1^{EM}/w_2^{EM} > \delta x_1$.

Lemma 1c. $\gamma > 1$ and $\gamma \geq \delta \Rightarrow w_1^{EM}/w_3^{EM} < \gamma x_2$.

Lemma 1d. $\gamma < 1$ and $\gamma \leq \delta \Rightarrow w_1^{EM}/w_3^{EM} > \gamma x_2$.

Lemma 1e. $\gamma, \delta > 1 \Rightarrow w_1^{EM}/w_i^{EM} > x_{i-1}$, $i = 4, \dots, n$.

Lemma 1f. $\gamma, \delta < 1 \Rightarrow w_1^{EM}/w_i^{EM} < x_{i-1}, i = 4, \dots, n.$

Lemma 1g. $\delta \leq \gamma \Leftrightarrow w_2^{EM}/w_3^{EM} \geq x_2/x_1.$

Lemma 1h. $\delta \geq 1 \Leftrightarrow w_2^{EM}/w_i^{EM} \leq x_{i-1}/x_1, i = 4, \dots, n.$

Lemma 1i. $\gamma \geq 1 \Leftrightarrow w_3^{EM}/w_i^{EM} \leq x_{i-1}/x_2, i = 4, \dots, n.$

Lemma 1j. $w_i^{EM}/w_j^{EM} = x_{j-1}/x_{i-1}, i, j = 4, \dots, n.$

The second group of lemmas correspond to Case 2A (γ and δ are in different rows, and matrix size is 4×4), i.e., the double perturbed PCM is written in form (4).

Lemma 2a. $\delta \geq 1 \Leftrightarrow w_1^{EM}/w_2^{EM} \leq \delta x_1.$

Lemma 2b. $\delta > 1, \gamma < 1 \Rightarrow w_1^{EM}/w_3^{EM} > x_2.$

Lemma 2c. $\delta < 1, \gamma > 1 \Rightarrow w_1^{EM}/w_3^{EM} < x_2.$

Lemma 2d. $\delta, \gamma < 1 \Rightarrow w_1^{EM}/w_4^{EM} < x_3.$

Lemma 2e. $\delta, \gamma > 1 \Rightarrow w_1^{EM}/w_4^{EM} > x_3.$

Lemma 2f. $\delta, \gamma < 1 \Rightarrow w_2^{EM}/w_3^{EM} > x_2/x_1.$

Lemma 2g. $\delta, \gamma > 1 \Rightarrow w_2^{EM}/w_3^{EM} < x_2/x_1.$

Lemma 2h. $\delta < 1, \gamma > 1 \Rightarrow w_2^{EM}/w_4^{EM} > x_3/x_1.$

Lemma 2i. $\delta > 1, \gamma < 1 \Rightarrow w_2^{EM}/w_4^{EM} < x_3/x_1.$

Lemma 2j. $\gamma \geq 1 \Leftrightarrow w_3^{EM}/w_4^{EM} \leq \gamma x_3/x_2.$

The last group of lemmas correspond to Case 2B, when γ and δ are in different rows, and matrix size is at least 5×5 , i.e., the double perturbed PCM is written in form (5).

Lemma 3a. $\gamma \geq 1 \Leftrightarrow w_3^{EM}/w_4^{EM} \leq \gamma x_3/x_2.$

Lemma 3b. $\delta \geq 1 \Leftrightarrow w_1^{EM}/w_2^{EM} \leq \delta x_1.$

Lemma 3c. $\delta \geq 1 \Leftrightarrow w_1^{EM}/w_i^{EM} \geq x_{i-1}, i = 5, \dots, n.$

Lemma 3d. $\delta \geq 1 \Leftrightarrow w_2^{EM}/w_i^{EM} \leq x_{i-1}/x_1, i = 5, \dots, n.$

Lemma 3e. $\gamma \geq 1 \Leftrightarrow w_3^{EM}/w_i^{EM} \geq x_{i-1}/x_2, i = 5, \dots, n.$

Lemma 3f.

$$\gamma > 1, \delta < 1 \Rightarrow w_2^{EM}/w_4^{EM} > x_3/x_1.$$

$$\gamma < 1, \delta > 1 \Rightarrow w_2^{EM}/w_4^{EM} < x_3/x_1.$$

Proof. Instead of the statement of the lemma, we will prove the following stronger statement:

$$\gamma \begin{smallmatrix} \geq \\ \leq \end{smallmatrix} \delta \Leftrightarrow w_2^{EM}/w_4^{EM} \begin{smallmatrix} \geq \\ \leq \end{smallmatrix} x_3/x_1.$$

Formula (21) is used in this proof.

$$\begin{aligned} & \frac{x_4}{x_1} \lambda (\gamma^2 - 2\gamma + \lambda^2 \gamma + 1) (1 + \delta \lambda - \delta) \\ & \begin{smallmatrix} \geq \\ \leq \end{smallmatrix} \frac{x_3}{x_1} \frac{x_4}{x_3} \lambda (\delta \lambda^2 + \lambda^3 \delta \gamma - \delta \gamma \lambda^2 - 2\lambda \delta \gamma - 2\delta + 2\delta \gamma - \gamma + 1 + \lambda \gamma + \delta^2 + \delta^2 \lambda \gamma - \delta^2 \gamma). \end{aligned}$$

This is further equivalent to

$$\begin{aligned} & \gamma^2 + \lambda \gamma^2 \delta - \gamma^2 \delta - 2\gamma - 2\lambda \gamma \delta + 2\gamma \delta + \lambda^2 \gamma + \lambda^3 \gamma \delta - \lambda^2 \gamma \delta + 1 + \lambda \delta - \delta \\ & \begin{smallmatrix} \geq \\ \leq \end{smallmatrix} \lambda^2 \delta + \lambda^3 \gamma \delta - \lambda^2 \gamma \delta - 2\lambda \gamma \delta - 2\delta + 2\gamma \delta - \gamma + 1 + \lambda \gamma + \delta^2 + \lambda \gamma \delta^2 - \gamma \delta^2. \end{aligned}$$

Further equivalent transformations yield

$$\begin{aligned} & \lambda^2 \gamma - \lambda^2 \delta + \lambda \delta - \lambda \gamma + \lambda \gamma^2 \delta - \lambda \gamma \delta^2 + \gamma^2 - \delta^2 + \gamma \delta^2 - \gamma^2 \delta + 2\delta - 2\gamma + \gamma - \delta \begin{smallmatrix} \geq \\ \leq \end{smallmatrix} 0 \\ & \lambda^2 (\gamma - \delta) + \lambda (\delta - \gamma) + \lambda \gamma \delta (\gamma - \delta) + (\gamma + \delta) (\gamma - \delta) + \gamma \delta (\delta - \gamma) + 2(\delta - \gamma) + (\gamma - \delta) \begin{smallmatrix} \geq \\ \leq \end{smallmatrix} 0 \\ & (\gamma - \delta) (\lambda^2 - \lambda + \lambda \gamma \delta + \gamma + \delta - \gamma \delta - 1) \begin{smallmatrix} \geq \\ \leq \end{smallmatrix} 0 \\ & (\gamma - \delta) (\lambda^2 - 2\lambda + \lambda \gamma \delta - \gamma \delta + \lambda - 1 + \gamma + \delta) \begin{smallmatrix} \geq \\ \leq \end{smallmatrix} 0 \\ & (\gamma - \delta) (\lambda (\lambda - 2) + \gamma \delta (\lambda - 1) + (\lambda - 1) + \gamma + \delta) \begin{smallmatrix} \geq \\ \leq \end{smallmatrix} 0. \end{aligned}$$

The second factor on the left hand side is always positive because $\lambda > n \geq 5$ and $\gamma, \delta > 0$. $\square$

Lemma 3g.

$$\begin{aligned} & \gamma, \delta > 1 \Rightarrow w_1^{EM}/w_4^{EM} > x_3. \\ & \gamma, \delta < 1 \Rightarrow w_1^{EM}/w_4^{EM} < x_3. \end{aligned}$$

Lemma 3h. $w_i^{EM}/w_j^{EM} = x_{j-1}/x_{i-1}$, $i, j = 5, \dots, n$.

Cases of $\delta = 1$ and $\gamma = 1$ are not covered by Lemmas 1a–3h due to Remark 2. Utilizing these lemmas, the main result of the paper can be proved:

Theorem 6. *The principal right eigenvector of a double perturbed PCM is efficient.*

Proof. As per Theorem 1, the strong connectedness of the digraph in Definition 2 needs to be shown. All possible digraphs are shown in Figures 2–4. The direction of each arc (where applicable) is determined by the corresponding lemma using Definition 2, which is labeled on the arc itself. In the cases where there is a node named i , this represents the complete subgraph of the rest of the nodes (consisting of $n - 3$ in Case 1 and $n - 4$ nodes in Case 2B). In these subgraphs there are bidirected arcs between any two nodes, due to Lemmas 1j and 3h. This is a strongly connected subgraph, and for any fixed $j \leq 3$ the direction of the arc between nodes i and j is the same for every $i \geq 4$ in Case 1 (see Lemmas 1e, 1f, 1h, 1i). Similarly, for any fixed $j \leq 4$ the

direction of the arc between nodes i and j is the same for every $i \geq 5$ in Case 2B (see Lemmas 3c, 3d, 3e). Hence, it can be contracted into a single node when analyzing strong connectedness. Figures 2, 3, 4 correspond to Cases 1, 2A, 2B respectively. For the strong connectedness of each digraph, it is sufficient to find a directed cycle. Unchecked arcs are denoted by dashed lines in Figures 2–4. Directed cycles are as follows:

Case 1 (Figure 2):

$$\begin{aligned}
&\delta > 1, \gamma > \delta : 1 \rightarrow i \rightarrow 2 \rightarrow 3 \rightarrow 1, \\
&\gamma > 1, \gamma < \delta : 1 \rightarrow i \rightarrow 3 \rightarrow 2 \rightarrow 1, \\
&\delta > 1, \gamma < 1 : 1 \rightarrow 3 \rightarrow i \rightarrow 2 \rightarrow 1, \\
&\delta < 1, \gamma < \delta : 1 \rightarrow 3 \rightarrow 2 \rightarrow i \rightarrow 1, \\
&\gamma < 1, \gamma > \delta : 1 \rightarrow 2 \rightarrow 3 \rightarrow i \rightarrow 1, \\
&\delta < 1, \gamma > 1 : 1 \rightarrow 2 \rightarrow i \rightarrow 3 \rightarrow 1.
\end{aligned}$$

Case 2A (Figure 3):

$$\begin{aligned}
&\delta > 1, \gamma > 1 : 1 \rightarrow 4 \rightarrow 3 \rightarrow 2 \rightarrow 1, \\
&\delta > 1, \gamma < 1 : 1 \rightarrow 3 \rightarrow 4 \rightarrow 2 \rightarrow 1, \\
&\delta < 1, \gamma < 1 : 1 \rightarrow 2 \rightarrow 3 \rightarrow 4 \rightarrow 1, \\
&\delta < 1, \gamma > 1 : 1 \rightarrow 2 \rightarrow 4 \rightarrow 3 \rightarrow 1.
\end{aligned}$$

Case 2B (Figure 4):

$$\begin{aligned}
&\delta > 1, \gamma > 1 : 1 \rightarrow 4 \rightarrow 3 \rightarrow i \rightarrow 2 \rightarrow 1, \\
&\delta > 1, \gamma < 1 : 1 \rightarrow i \rightarrow 3 \rightarrow 4 \rightarrow 2 \rightarrow 1, \\
&\delta < 1, \gamma < 1 : 1 \rightarrow 2 \rightarrow i \rightarrow 3 \rightarrow 4 \rightarrow 1, \\
&\delta < 1, \gamma > 1 : 1 \rightarrow 2 \rightarrow 4 \rightarrow 3 \rightarrow i \rightarrow 1.
\end{aligned}$$

The presence of a directed cycle implies strong connectedness for all of the digraphs, which implies efficiency in all cases by Theorem 1. $\square$

Figure 2: The digraph of the principal right eigenvector in Case 1 is strongly connected, independently of the orientation of dashed arcs that have not been analyzed




Figure 3: The digraph of the principal right eigenvector in Case 2A is strongly connected, independently of the orientation of dashed arcs that have not been analyzed





Figure 4: The digraph of the principal right eigenvector in Case 2B is strongly connected, independently of the orientation of dashed arcs that have not been analyzed



4 Conclusions

In the paper we used linear algebraic methods to derive explicit formulas for the principal eigenvector of double perturbed PCM. We also used a necessary and sufficient condition for efficiency which uses a directed graph representation (the weight vector is efficient if and only if this graph is strongly connected) developed by Blanquero, Carrizosa and Conde [4]. Double perturbed PCMs had to be divided into three cases in order to get explicit formulas for every case. In all three cases the digraph has been studied arc by arc, however not all arcs had to be studied in order to determine strong connectedness. Utilizing all these tools, we have shown in the paper, that the often used eigenvector method produces an efficient weight vector in the case of double perturbed PCMs. This is an extension of our earlier result for simple perturbed PCMs [1]. Extension to triple perturbed and further is not possible, since all PCMs of at least 4×4 size which are not (at most) double perturbed are triple perturbed, and there are examples of inefficiency at size 4×4 . Thus, while in some cases (e.g. when all perturbed elements are in different rows/columns) it may be possible to show efficiency, for all triple perturbed PCMs this is impossible. Furthermore, a triple perturbed PCM can be equivalent to five separate basic cases (see [7, Fig. 7]), which may need to be further divided into more subcases, making the efficiency analysis of triple perturbed PCMs difficult. A full characterization of the efficiency of the principal right eigenvector is still an open question, and a possible subject of future research.

References

- [1] K. Ábele-Nagy and S. Bozóki. Efficiency analysis of simple perturbed pairwise comparison matrices. *Fundamenta Informaticae*, *accepted*, 2016. <http://arxiv.org/abs/1505.06849>.
- [2] M. Anholcer and J. Fülöp. Deriving priorities from inconsistent PCM using the network algorithms. *manuscript*, 2015. <http://arxiv.org/pdf/1510.04315>.
- [3] G. Bajwa, E. U. Choo, and W. C. Wedley. Effectiveness analysis of deriving priority vectors from reciprocal pairwise comparison matrices. *Asia-Pacific Journal of Operational Research*, 25(3):279–299, 2008.
- [4] R. Blanquero, E. Carrizosa, and E. Conde. Inferring efficient weights from pairwise comparison matrices. *Mathematical Methods of Operations Research*, 64(2):271–284, 2006.
- [5] S. Bozóki. Inefficient weights from pairwise comparison matrices with arbitrarily small inconsistency. *Optimization*, 63(12):1893–1901, 2014.
- [6] S. Bozóki and J. Fülöp. Efficient weight vectors from pairwise comparison matrices. *manuscript*, 2015. <http://arxiv.org/abs/1602.03311>.
- [7] S. Bozóki, J. Fülöp, and A. Poesz. On pairwise comparison matrices that can be made consistent by the modification of a few elements. *Central European Journal of Operations Research*, 19(2):157–175, 2011.
- [8] M. Brunelli and M. Fedrizzi. Axiomatic properties of inconsistency indices for pairwise comparisons. *Journal of the Operational Research Society*, 66:1–15, 2014.
- [9] E. U. Choo and W. C. Wedley. A common framework for deriving preference values from pairwise comparison matrices. *Computers & Operations Research*, 31(6):893–908, 2004.
- [10] E. Conde and M. d. l. P. R. Pérez. A linear optimization problem to derive relative weights using an interval judgement matrix. *European Journal of Operational Research*, 201(2):537–544, 2010.
- [11] W. D. Cook and M. Kress. Deriving weights from pairwise comparison ratio matrices: An axiomatic approach. *European Journal of Operational Research*, 37(3):355–362, 1988.
- [12] T. K. Dijkstra. On the extraction of weights from pairwise comparison matrices. *Central European Journal of Operations Research*, 21(1):103–123, 2013.
- [13] M. Ehrgott. *Multicriteria Optimization*. volume 491 of *Lecture Notes in Economics and Mathematical Systems*. Springer Verlag, Berlin, 2000.
- [14] A. Farkas. The analysis of the principal eigenvector of pairwise comparison matrices. *Acta Polytechnica Hungarica*, 4(2):99–115, 2007.

- [15] A. Farkas, P. Rózsa, and E. Stubnya. Transitive matrices and their applications. *Linear Algebra and its Applications*, 302–303:423–433, 1999.
- [16] M. Fedrizzi. Obtaining non-dominated weights from preference relations through norm-induced distances. *XXXVII Meeting of the Italian Association for Mathematics Applied to Economic and Social Sciences (AMASES), September 5–7, 2013, Stresa, Italy*.
- [17] B. Golany and M. Kress. A multicriteria evaluation of methods for obtaining weights from ratio-scale matrices. *European Journal of Operational Research*, 69(2):210–220, 1993.
- [18] D. A. Harville. *Matrix Algebra From a Statistician's Perspective*. Springer, 2008.
- [19] A. Poesz. Empirical pairwise comparison matrices (epcm) – an on-line collection from real decisions, version epcm-october-2009, 2009.
- [20] T. L. Saaty. A scaling method for priorities in hierarchical structures. *Journal of Mathematical Psychology*, 15(3):234–281, 1977.
- [21] J. Sherman and W. J. Morrison. Adjustment of an inverse matrix corresponding to a change in one element of a given matrix. *Annals of Mathematical Statistics*, 21(1):124–127, 1950.
- [22] R. E. Steuer. *Multiple Criteria Optimization: Theory, Computation, and Application*. Wiley Series in Probability and Mathematical Statistics. Wiley, 1986.

Appendix: Proofs of Lemmas 1a–3h

Remark 4. *Formulas (9)–(21) are positive.*

Proof. It is sufficient to prove the positivity of any arbitrary element of each formula, because the Perron–Frobenius theorem then guarantees the positivity for the vectors as well. The conclusions of the proofs generally follow from $x_i > 0 \forall i = 1, \dots, n$, $\gamma, \delta > 0$ and $\lambda > n \geq 4$ (or $n \geq 5$ in Case 2B). The proof for each formula follows:

Formula (9): Positivity is apparent for w_1^{EM} .

Formula (10):

$$\begin{aligned} w_1^{EM} &= x_1 \gamma \lambda [\delta \lambda - (n-2)\delta + \gamma + n - 3] \\ &= x_1 \gamma \lambda [\delta(\lambda - n + 2) + \gamma + (n-3)]. \end{aligned}$$

Formula (11):

$$\begin{aligned} w_1^{EM} &= x_2 \delta \lambda [\delta + \gamma \lambda - (n-2)\gamma + n - 3] \\ &= x_2 \delta \lambda [\delta + \gamma(\lambda - n + 2) + (n-3)]. \end{aligned}$$

Formula (12): Positivity is apparent for w_1^{EM} .

Formula (13):

$$w_2^{EM} = \frac{1}{x_1} [\lambda^2 \gamma - 2\lambda \gamma + \delta + 2\lambda \delta \gamma - 2\delta \gamma + \delta \gamma^2]$$

$$= \frac{1}{x_1} [\lambda\gamma(\lambda - 2) + \delta + 2\delta\gamma(\lambda - 1) + \delta\gamma^2].$$

Formula (14):

$$\begin{aligned} w_1^{EM} &= x_1[\delta\gamma\lambda^2 - 2\lambda\delta\gamma + 1 + 2\lambda\gamma - 2\gamma + \gamma^2] \\ &= x_1[\delta\gamma\lambda(\lambda - 2) + 1 + 2\gamma(\lambda - 1) + \gamma^2]. \end{aligned}$$

Formula (15):

$$\begin{aligned} w_1^{EM} &= x_2\delta(1 + \lambda\gamma - \gamma)(\delta + \lambda - 1) \\ &= x_2\delta[1 + \gamma(\lambda - 1)][\delta + (\lambda - 1)]. \end{aligned}$$

Formula (16):

$$\begin{aligned} w_3^{EM} &= \frac{x_3}{x_2} [2\lambda\delta + \delta\gamma\lambda^2 - 2\lambda\delta\gamma - 2\delta + 1 + \delta^2] \\ &= \frac{x_3}{x_2} [2\delta(\lambda - 1) + \delta\gamma\lambda(\lambda - 2) + 1 + \delta^2]. \end{aligned}$$

From here on in the proof, $n \geq 5$.

Formula (17): w_3^{EM} in formula (17) is the same as λw_3^{EM} in formula (13), which is already proven to be positive.

Formula (18): w_3^{EM} in formula (18) is the same as λw_3^{EM} in formula (14), which is already proven to be positive.

Formula (19):

$$\begin{aligned} w_1^{EM} &= x_2\delta\lambda(1 + \lambda\gamma - \gamma)(\delta + \lambda - 1) \\ &= x_2\delta\lambda[1 + \gamma(\lambda - 1)][\delta + (\lambda - 1)]. \end{aligned}$$

Formula (20):

$$\begin{aligned} w_2^{EM} &= \frac{x_3}{x_1}\lambda(\gamma + \lambda - 1)(1 + \delta\lambda - \delta) \\ &= \frac{x_3}{x_1}\lambda[\gamma + (\lambda - 1)][1 + \delta(\lambda - 1)]. \end{aligned}$$

Formula (21):

$$\begin{aligned} w_1^{EM} &= x_4\delta\lambda(\gamma^2 - 2\gamma + \lambda^2\gamma + 1)(\delta + \lambda - 1) \\ &= x_4\delta\lambda[\gamma^2 + \gamma(\lambda^2 - 2) + 1][\delta + (\lambda - 1)]. \end{aligned}$$

□

Lemma 1a (Case 1). $\delta > 1$ and $\delta \geq \gamma \Rightarrow w_1^{EM}/w_2^{EM} < \delta x_1$.

Proof. Using formula (10),

$$\frac{w_1^{EM}}{w_2^{EM}} = x_1 \frac{\gamma\lambda(\delta\lambda - (n-2)\delta + \gamma + n-3)}{\gamma\lambda^3 - (n-1)\gamma\lambda^2 - (n-3)(\gamma^2 - 2\gamma + 1)}.$$

Substitute $\lambda = \lambda_{\max}$ in the characteristic polynomial $p_{\mathbf{P}}(\lambda)$ by Theorem 3:

$$(-1)^n \lambda^{n-3} \left(\lambda^3 - n\lambda^2 - \left(\frac{\gamma}{\delta} + \frac{\delta}{\gamma} \right) - (n-3) \left(\gamma + \delta + \frac{1}{\gamma} + \frac{1}{\delta} \right) + 4n - 10 \right) = 0,$$

which can be transformed to

$$\gamma\delta\lambda^3 - \gamma\delta n\lambda^2 = \gamma^2 + \delta^2 + (n-3)(\gamma^2\delta + \gamma\delta^2 + \delta + \gamma) - \gamma\delta(4n-10). \quad (22)$$

The statement to be proven is equivalent to

$$\gamma\lambda(\delta\lambda + \gamma - (n-2)\delta + n-3) < \delta(\gamma\lambda^3 - (n-1)\gamma\lambda^2 - (n-3)(\gamma-1)^2).$$

Using (22) this is further equivalent to

$$(\gamma\delta\lambda^3 - \gamma\delta n\lambda^2) + \lambda(\gamma\delta(n-2) - \gamma^2 - \gamma n + 3\gamma) - \delta(n-3)(\gamma-1)^2 > 0.$$

Now apply further equivalent transformations:

$$\begin{aligned} \gamma\lambda(\delta(n-2) - \gamma - n + 3) + \gamma^2 + \delta^2 - (n-3)(\delta\gamma^2 - 2\delta\gamma + \delta) \\ + (n-3)(\delta^2\gamma + \delta\gamma^2 + \delta + \gamma) - \delta\gamma(4n-10) > 0 \end{aligned}$$

$$\begin{aligned} \gamma\lambda((\delta-1)(n-3) + \delta - \gamma) + \gamma^2 + \delta^2 \\ + (n-3)(\delta^2\gamma + 2\delta\gamma + \gamma) - 4\delta\gamma(n-3) - 2\delta\gamma > 0 \end{aligned}$$

$$\gamma\lambda((\delta-1)(n-3) + (\delta-\gamma)) + \gamma(n-3)(\delta-1)^2 + (\delta-\gamma)^2 > 0.$$

□

Lemma 1b (Case 1). $\delta < 1$ and $\delta \leq \gamma \Rightarrow w_1^{EM}/w_2^{EM} > \delta x_1$.

Proof. According to formula (10)

$$\frac{w_1^{EM}}{w_2^{EM}} = x_1 \frac{\gamma\lambda(\delta\lambda - (n-2)\delta + \gamma + n-3)}{\gamma\lambda^3 - (n-1)\gamma\lambda^2 - (n-3)(\gamma^2 - 2\gamma + 1)}.$$

Transforming (22) similar to Lemma 1a,

$$\gamma\lambda((\delta-1)(n-3) + \delta - \gamma) + \gamma(n-3)(\delta-1)^2 + (\delta-\gamma)^2 < 0.$$

Transforming this further yields

$$\begin{aligned} \gamma(\delta-1)(n-3)(\lambda + (\delta-1)) + \gamma\lambda(\delta-\gamma) + (\delta-\gamma)^2 < 0 \\ \gamma(\delta-1)(n-3)(\lambda + (\delta-1)) + (\delta-\gamma)(\gamma(\lambda-1) + \delta) < 0. \end{aligned}$$

□

Lemma 1c (Case 1). $\gamma > 1$ and $\gamma \geq \delta \Rightarrow w_1^{EM}/w_3^{EM} < \gamma x_2$.

Proof. The proof follows from switching the role of δ and γ in the proof of Lemma 1a. □

Lemma 1d (Case 1). $\gamma < 1$ and $\gamma \leq \delta \Rightarrow w_1^{EM}/w_3^{EM} > \gamma x_2$.

Proof. The proof follows from switching the role of δ and γ in the proof of Lemma 1b. $\square$

Lemma 1e (Case 1). $\gamma, \delta > 1 \Rightarrow w_1^{EM}/w_i^{EM} > x_{i-1}, i = 4, \dots, n.$

Proof. According to formula (9)

$$\frac{w_1^{EM}}{w_i^{EM}} = x_{i-1} \frac{\gamma\delta\lambda(\lambda - n + 1)}{\gamma + \delta + \gamma\delta\lambda - 2\gamma\delta},$$

which means the statement to be proven is equivalent to

$$\gamma\delta\lambda(\lambda - n + 1) > \gamma + \delta + \gamma\delta\lambda - 2\gamma\delta.$$

Further equivalent transformations yield

$$\begin{aligned} (\gamma\delta\lambda(\lambda - n)) + (2\gamma\delta - \gamma - \delta) &> 0 \\ \gamma\delta\lambda(\lambda - n) + (\delta - 1)(\gamma - 1) + (\delta\gamma - 1) &> 0. \end{aligned}$$

$\square$

Lemma 1f (Case 1). $\gamma, \delta < 1 \Rightarrow w_1^{EM}/w_i^{EM} < x_{i-1}, i = 4, \dots, n.$

Proof. According to formula (12)

$$\frac{w_1^{EM}}{w_i^{EM}} = x_{i-1} \frac{\gamma\delta\lambda(\delta + \gamma + \lambda - 2)}{\gamma\delta\lambda^2 - 4\gamma\delta + \gamma + \delta + \delta^2\gamma + \gamma^2\delta}.$$

Applying further equivalent transformations

$$\begin{aligned} \frac{\gamma\delta\lambda(\delta + \gamma + \lambda - 2)}{\gamma\delta\lambda^2 - 4\gamma\delta + \gamma + \delta + \delta^2\gamma + \gamma^2\delta} &< 1 \\ \gamma\delta\lambda(\delta + \gamma + \lambda - 2) &< \gamma\delta(\lambda^2 - 4) + \gamma + \delta + \delta^2\gamma + \gamma^2\delta \\ \lambda(\delta + \gamma + \lambda - 2) &< \lambda^2 - 4 + \frac{1}{\delta} + \frac{1}{\gamma} + \delta + \gamma \\ 0 &< \lambda^2 - 4 + \frac{1}{\delta} + \frac{1}{\gamma} + \delta + \gamma - \lambda\delta - \lambda\gamma - \lambda^2 + 2\lambda \\ 0 &< 2(\lambda - 2) + (1 - \lambda)(\delta + \gamma) + \frac{1}{\delta} + \frac{1}{\gamma} \\ 0 &< 2(\lambda - 1) - 2 + (1 - \lambda)(\delta + \gamma) + \frac{1}{\delta} + \frac{1}{\gamma} \\ 0 &< (\lambda - 1)(2 - \delta - \gamma) + \frac{1}{\delta} + \frac{1}{\gamma} - 2. \end{aligned}$$

$\square$

Lemma 1g (Case 1). $\delta \lessgtr \gamma \Leftrightarrow w_2^{EM}/w_3^{EM} \lessgtr x_2/x_1.$

Proof. According to formula (9), we need to consider

$$\frac{w_2^{EM}}{w_3^{EM}} = \frac{x_2}{x_1} \cdot \frac{\gamma\lambda - (n-2)\gamma + \delta + (n-3)\gamma\delta}{\delta\lambda - (n-2)\delta + \gamma + (n-3)\gamma\delta} \geq 1.$$

Applying further equivalent transformations

$$\begin{aligned} \gamma\lambda - (n-2)\gamma + \delta + (n-3)\gamma\delta &\geq \delta\lambda - (n-2)\delta + \gamma + (n-3)\gamma\delta \\ \lambda(\gamma - \delta) - (n-2)(\gamma - \delta) + \delta - \gamma &\geq 0 \\ (\gamma - \delta)(\lambda - n + 1) &\geq 0. \end{aligned}$$

The third factor is positive because $\lambda > n$. □

Lemma 1h (Case 1). $\delta \geq 1 \Leftrightarrow w_2^{EM}/w_i^{EM} \leq x_{i-1}/x_1, i = 4, \dots, n$.

Proof. According to formula (9)

$$\frac{w_2^{EM}}{w_i^{EM}} = \frac{x_{i-1}}{x_1} \cdot \frac{\gamma\lambda - (n-2)\gamma + \delta + (n-3)\gamma\delta}{\gamma + \delta + \gamma\delta\lambda - 2\gamma\delta}.$$

Equivalent transformations yield

$$\begin{aligned} \gamma\lambda - (n-2)\gamma + \delta + (n-3)\gamma\delta &< \gamma + \delta + \gamma\delta\lambda - 2\gamma\delta \\ 0 &< \gamma(\delta - 1)(\lambda - n + 1). \end{aligned}$$

The third factor is positive because $\lambda > n$. □

Lemma 1i (Case 1). $\gamma \geq 1 \Leftrightarrow w_3^{EM}/w_i^{EM} \leq x_{i-1}/x_2, i = 4, \dots, n$.

Proof. The proof follows from switching the role of δ and γ in the proof of Lemma 1h. □

Lemma 1j (Case 1). $w_i^{EM}/w_j^{EM} = x_{j-1}/x_{i-1}, i, j = 4, \dots, n$.

Proof. It follows from each of formulas (9)–(12). □

Lemma 2a (Case 2A). $\delta \geq 1 \Leftrightarrow w_1^{EM}/w_2^{EM} \leq \delta x_1$.

Proof. Formula (16) is used for this proof. Multiplying both sides by w_2^{EM} , the statement to be proven can be written as:

$$\begin{aligned} x_3\delta(\lambda\gamma + \lambda^2 - 2\lambda - \gamma + 1 + \lambda\delta - \delta + \delta\gamma) \\ \leq \delta x_1 \frac{x_3}{x_1} (\gamma + \lambda - 1 + \delta\lambda^2 - 2\lambda\delta + \delta + \lambda\delta\gamma - \delta\gamma). \end{aligned} \quad (23)$$

Further equivalent transformations yield:

$$\begin{aligned} 0 &\leq \lambda^2\delta - \lambda^2 + 3\lambda - \lambda\gamma - 3\lambda\delta + \lambda\delta\gamma - 2\delta\gamma + 2\gamma + 2\delta - 2 \\ 0 &\leq \lambda^2(\delta - 1) + \lambda\gamma(\delta - 1) + 3\lambda(1 - \delta) + 2\gamma(1 - \delta) + 2(\delta - 1) \\ 0 &\leq (\delta - 1)(\lambda(\lambda - 3) + \gamma(\lambda - 2) + 2). \end{aligned}$$

The second factor on the right hand side is always positive because $\lambda > n = 4$ and $\gamma, \delta > 0$. □

Lemma 2b (Case 2A). $\delta > 1, \gamma < 1 \Rightarrow w_1^{EM}/w_3^{EM} > x_2$.

Proof. Formula (14) is used in this proof. Multiplying both sides by w_3^{EM} , the statement of the lemma is equivalent to:

$$\begin{aligned} x_1(\delta\gamma\lambda^2 - 2\lambda\delta\gamma + 1 + 2\lambda\gamma - 2\gamma + \gamma^2) \\ < x_2 \frac{x_1}{x_2} \gamma (\lambda\gamma + \lambda^2 - 2\lambda - \gamma + 1 + \lambda\delta - \delta + \delta\gamma). \end{aligned}$$

Further equivalent transformations yield:

$$\begin{aligned} 0 &< \lambda^2\gamma - \lambda^2\gamma\delta - 4\lambda\gamma + \lambda\gamma^2 + 3\lambda\delta\gamma - 2\gamma^2 + 3\gamma + \delta\gamma^2 - \delta\gamma - 1 \\ 0 &< \lambda^2\gamma(1 - \delta) + \lambda\gamma(\gamma - 1) + 3\lambda\gamma(\delta - 1) + (\gamma - 1) + 2\gamma(1 - \gamma) + \delta\gamma(\gamma - 1) \\ 0 &< (1 - \delta)\lambda\gamma(\lambda - 3) + (\gamma - 1)(\gamma(\lambda - 2) + \delta\gamma + 1). \end{aligned} \quad (24)$$

The second factor on the right hand side is always positive because $\lambda > n = 4$ and $\gamma, \delta > 0$. $\square$

Lemma 2c (Case 2A). $\delta < 1, \gamma > 1 \Rightarrow w_1^{EM}/w_3^{EM} < x_2$.

Proof. The proof follows from the right hand side of (24) being positive in the case of $\delta < 1, \gamma > 1$. $\square$

Lemma 2d (Case 2A). $\delta, \gamma < 1 \Rightarrow w_1^{EM}/w_4^{EM} < x_3$.

Proof. Again, formula (14) is used for this proof. Multiplying both sides by w_4^{EM} , the statement to be proven is equivalent to:

$$\begin{aligned} x_1(\delta\gamma\lambda^2 - 2\lambda\delta\gamma + 1 + 2\lambda\gamma - 2\gamma + \gamma^2) \\ < x_3 \frac{x_1}{x_3} (\lambda + \lambda^2\gamma - 2\lambda\gamma - 1 + \gamma + \delta + \lambda\delta\gamma - \delta\gamma). \end{aligned}$$

Further equivalent transformations yield:

$$\begin{aligned} \lambda^2\gamma\delta - \lambda^2\gamma + 4\lambda\gamma - 3\lambda\delta\gamma - \lambda + \gamma^2 - 3\gamma + \delta\gamma - \delta + 2 &< 0 \\ (\delta - 1)(\lambda^2\gamma - 3\lambda\gamma) + (\gamma - 1)(\lambda + \gamma - 2 + \delta) &< 0 \\ (\delta - 1)\lambda\gamma(\lambda - 3) + (\gamma - 1)((\lambda - 2) + \gamma + \delta) &< 0. \end{aligned} \quad (25)$$

The left hand side is negative if $\gamma, \delta < 1$, because $\lambda > n = 4$. $\square$

Lemma 2e (Case 2A). $\delta, \gamma > 1 \Rightarrow w_1^{EM}/w_4^{EM} > x_3$.

Proof. The proof follows from the left hand side of (25) being positive if $\gamma, \delta > 1$. $\square$

Lemma 2f (Case 2A). $\delta, \gamma < 1 \Rightarrow w_2^{EM}/w_3^{EM} > x_2/x_1$.

Proof. Formula (13) is used in this proof. Multiplying both sides by w_3^{EM} , the statement of the lemma can be written as:

$$\frac{1}{x_1} (\lambda^2\gamma - 2\lambda\gamma + \delta + 2\lambda\delta\gamma - 2\delta\gamma + \delta\gamma^2)$$

$$> \frac{x_2}{x_1} \frac{1}{x_2} \gamma (\gamma + \lambda - 1 + \delta \lambda^2 - 2\lambda\delta + \delta + \lambda\delta\gamma - \delta\gamma).$$

Further equivalent transformations yield:

$$\begin{aligned} 0 &> \lambda^2\gamma\delta - \lambda^2\gamma - 4\lambda\delta\gamma + 3\lambda\gamma + \lambda\delta\gamma^2 + \gamma^2 - 2\delta\gamma^2 + 3\delta\gamma - \gamma - \delta \\ 0 &> (\delta - 1)(\lambda^2\gamma - 3\lambda\gamma) + (\gamma - 1)(\lambda\delta\gamma - 2\delta\gamma + \delta + \gamma) \\ 0 &> (\delta - 1)\lambda\gamma(\lambda - 3) + (\gamma - 1)(\delta\gamma(\lambda - 2) + \delta + \gamma). \end{aligned} \quad (26)$$

The right hand side is negative if $\delta, \gamma < 1$, because $\lambda > n = 4$. $\square$

Lemma 2g (Case 2A). $\delta, \gamma > 1 \Rightarrow w_2^{EM}/w_3^{EM} < x_2/x_1$.

Proof. The proof follows from the right hand side of (26) being positive if $\delta, \gamma > 1$. $\square$

Lemma 2h (Case 2A). $\delta < 1, \gamma > 1 \Rightarrow w_2^{EM}/w_4^{EM} > x_3/x_1$.

Proof. Again, formula (13) is used in this proof. Multiplying both sides by w_4^{EM} , the statement to be proven is equivalent to:

$$\begin{aligned} \frac{1}{x_1} (\lambda^2\gamma - 2\lambda\gamma + \delta + 2\lambda\delta\gamma - 2\delta\gamma + \delta\gamma^2) \\ > \frac{x_3}{x_1} \frac{1}{x_3} (1 + \lambda\gamma - \gamma + \lambda\delta - \delta + \delta\gamma\lambda^2 - 2\lambda\delta\gamma + \delta\gamma). \end{aligned}$$

Further equivalent transformations yield:

$$\begin{aligned} 0 &> \lambda^2\gamma\delta - \lambda^2\gamma - 4\lambda\delta\gamma + 3\lambda\gamma + \lambda\delta + 3\delta\gamma - \delta\gamma^2 - 2\delta - \gamma + 1 \\ 0 &> (\delta - 1)(\lambda^2\gamma - 3\lambda\gamma) + (1 - \gamma)(\lambda\delta - 2\delta + \delta\gamma + 1) \\ 0 &> (\delta - 1)\lambda\gamma(\lambda - 3) + (1 - \gamma)(\delta(\lambda - 2) + \delta\gamma + 1). \end{aligned} \quad (27)$$

The right hand side of (27) is negative, if $\delta < 1, \gamma > 1$, because $\lambda > n = 4$. $\square$

Lemma 2i (Case 2A). $\delta > 1, \gamma < 1 \Rightarrow w_2^{EM}/w_4^{EM} < x_3/x_1$.

Proof. The proof follows from the right hand side of (27) being positive if $\delta > 1, \gamma < 1$. $\square$

Lemma 2j (Case 2A). $\gamma \gtrless 1 \Leftrightarrow w_3^{EM}/w_4^{EM} \lesseqgtr \gamma x_3/x_2$.

Proof. Once again, formula (13) is used for the proof. Multiplying both sides by w_4^{EM} , the first statement (for $\gamma > 1$) becomes equivalent to:

$$\begin{aligned} \frac{1}{x_2} \gamma (\gamma + \lambda - 1 + \delta \lambda^2 - 2\lambda\delta + \delta + \lambda\delta\gamma - \delta\gamma) \\ \lesseqgtr \gamma \frac{x_3}{x_2} \frac{1}{x_3} (1 + \lambda\gamma - \gamma + \lambda\delta - \delta + \delta\gamma\lambda^2 - 2\lambda\delta\gamma + \delta\gamma). \end{aligned} \quad (28)$$

Applying further equivalent transformations:

$$\begin{aligned} 0 &\lesseqgtr \lambda^2\delta\gamma - \lambda^2\delta - 3\lambda\delta\gamma + 3\lambda\delta + \lambda\gamma - \lambda + 2\delta\gamma - 2\delta - 2\gamma + 2 \\ 0 &\lesseqgtr (\gamma - 1)(\lambda^2\delta - 3\lambda\delta + \lambda + 2\delta - 2) \\ 0 &\lesseqgtr (\gamma - 1)(\lambda\delta(\lambda - 3) + (\lambda - 2) + 2\delta). \end{aligned} \quad (29)$$

The second factor on the right hand side of (29) is positive because $\lambda > n = 4$ and $\gamma, \delta > 0$. $\square$

Lemma 3a (Case 2B). $\gamma \geq 1 \Leftrightarrow w_3^{EM}/w_4^{EM} \leq \gamma x_3/x_2$.

Proof. Using formula (17) the proof is similar to the proof of Lemma 2j, the only difference is in (28) where both sides are multiplied by λ , which immediately cancel each other. $\square$

Lemma 3b (Case 2B). $\delta \geq 1 \Leftrightarrow w_1^{EM}/w_2^{EM} \leq \delta x_1$.

Proof. Using formula (21) the proof is similar to the proof of Lemma 2a, the only difference is in (23) where both sides (the formula for w_1^{EM} and w_2^{EM}) are multiplied by λ , which immediately cancel each other. This may not be apparent about w_2^{EM} , but

$$(\gamma + \lambda - 1)(1 + \delta\lambda - \delta) = \gamma + \lambda - 1 + \lambda\gamma\delta + \lambda^2\delta - \lambda\delta - \gamma\delta - \lambda\delta + \delta$$

which, after reduction, gives the same formula. $\square$

Lemma 3c (Case 2B). $\delta \geq 1 \Leftrightarrow w_1^{EM}/w_i^{EM} \geq x_{i-1}$, $i = 5, \dots, n$.

Proof. Formula (19) is used for this proof.

$$\begin{aligned} x_2\delta\lambda(1 + \lambda\gamma - \gamma)(\delta + \lambda - 1) &\geq x_{i-1}\frac{x_2}{x_{i-1}}(1 + \lambda\gamma - \gamma)(\delta\lambda^2 + 1 - 2\delta + \delta^2) \\ \lambda\delta^2 + \lambda^2\delta - \lambda\delta &\geq \delta\lambda^2 + 1 - 2\delta + \delta^2 \\ \lambda\delta(\delta - 1) + \delta(1 - \delta) + (\delta - 1) &\geq 0 \\ (\delta - 1)(\delta(\lambda - 1) + 1) &\geq 0. \end{aligned}$$

The second factor on the left hand side is always positive because $\lambda > n \geq 5$ and $\delta > 0$. $\square$

Lemma 3d (Case 2B). $\delta \geq 1 \Leftrightarrow w_2^{EM}/w_i^{EM} \leq x_{i-1}/x_1$, $i = 5, \dots, n$.

Proof. Again, formula (19) is used in the proof.

$$\begin{aligned} \frac{x_2}{x_1}\lambda(1 + \lambda\gamma - \gamma)(1 + \delta\lambda - \delta) &\leq \frac{x_{i-1}}{x_1}\frac{x_2}{x_{i-1}}(1 + \lambda\gamma - \gamma)(\delta\lambda^2 + 1 - 2\delta + \delta^2) \\ \lambda + \lambda^2\delta - \delta\lambda &\leq \delta\lambda^2 + 1 - 2\delta + \delta^2 \\ 0 &\leq \lambda\delta - \lambda + \delta^2 - 2\delta + 1 \\ 0 &\leq \lambda(\delta - 1) + (\delta - 1)^2 \\ 0 &\leq (\delta - 1)((\lambda - 1) + \delta). \end{aligned}$$

The second factor on the right hand side is always positive because $\lambda > n \geq 5$ and $\delta > 0$. $\square$

Lemma 3e (Case 2B). $\gamma \geq 1 \Leftrightarrow w_3^{EM}/w_i^{EM} \geq x_{i-1}/x_2$, $i = 5, \dots, n$.

Proof. Formula (18) is used in this proof.

$$\begin{aligned} \frac{x_1}{x_2}\gamma\lambda(\lambda\gamma + \lambda^2 - 2\lambda - \gamma + 1 + \delta\lambda - \delta + \delta\gamma) \\ \geq \frac{x_4}{x_2}\frac{x_1}{x_4}(\lambda\gamma^2 - 2\lambda\gamma + \lambda^3\gamma + \lambda - \gamma^2 + 2\gamma - \lambda^2\gamma - 1 + \delta - 2\delta\gamma + \delta\gamma^2 + \delta\gamma\lambda^2). \end{aligned}$$

Further equivalent transformations yield:

$$\begin{aligned}\lambda^2\gamma^2 - \lambda^2\gamma - 2\lambda\gamma^2 + 3\lambda\gamma + \lambda\gamma^2\delta - \lambda\delta\gamma - \lambda + \gamma^2 - 2\gamma + 1 - \delta + 2\delta\gamma - \delta\gamma^2 &\geq 0 \\ (\gamma - 1)(\lambda^2\gamma - 2\lambda\gamma + \lambda + \lambda\delta\gamma + (\gamma - 1) - \delta(\gamma - 1)) &\geq 0 \\ (\gamma - 1)(\lambda\gamma(\lambda - 2) + (\lambda - 1) + \delta\gamma(\lambda - 1) + \gamma + \delta) &\geq 0.\end{aligned}$$

The second factor on the left hand side is always positive because $\lambda > n \geq 5$ and $\gamma, \delta > 0$. $\square$

Lemma 3g (Case 2B).

$$\gamma, \delta > 1 \Rightarrow w_1^{EM}/w_4^{EM} > x_3.$$

$$\gamma, \delta < 1 \Rightarrow w_1^{EM}/w_4^{EM} < x_3.$$

Proof. Instead of the above statement, we will prove the following stronger statement:

$$\gamma\delta \geq 1 \Leftrightarrow w_1^{EM}/w_4^{EM} \geq x_3.$$

Formula (21) is used in this proof.

$$\begin{aligned}&x_4\delta\lambda(\gamma^2 - 2\gamma + \lambda^2\gamma + 1)(\delta + \lambda - 1) \\ &\geq x_3\frac{x_4}{x_3}\lambda(\delta\lambda^2 + \lambda^3\delta\gamma - \delta\gamma\lambda^2 - 2\lambda\delta\gamma - 2\delta + 2\delta\gamma - \gamma + 1 + \lambda\gamma + \delta^2 + \delta^2\lambda\gamma - \delta^2\gamma).\end{aligned}$$

Further equivalent transformations yield:

$$\begin{aligned}\lambda^2\delta^2\gamma - \lambda^2\delta + \lambda\gamma^2\delta - \lambda\gamma\delta^2 + \lambda\delta - \lambda\gamma + \gamma^2\delta^2 - \gamma\delta^2 - \delta\gamma^2 + \delta + \gamma - 1 &\geq 0 \\ (\delta\gamma - 1)(\lambda^2\delta + \lambda\gamma - \lambda\delta + \delta\gamma + 1 - \delta - \gamma) &\geq 0 \\ (\delta\gamma - 1)(\lambda\delta(\lambda - 2) + \gamma(\lambda - 1) + \delta(\lambda - 1) + \delta\gamma + 1) &\geq 0.\end{aligned}$$

The second factor is always positive because $\lambda > n \geq 5$ and $\gamma, \delta > 0$. The first factor is positive exactly if $\gamma\delta > 1$, and negative exactly if $\gamma\delta < 1$. $\square$

Lemma 3h (Case 2B). $w_i^{EM}/w_j^{EM} = x_{j-1}/x_{i-1}$, $i, j = 5, \dots, n$.

Proof. It follows from each of formulas (17)–(21). $\square$